

**Table 1.11A Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2010 and 2011**

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
<b>TOTAL</b>	119,933	121,078	38,908	38,287	22,648	22,454
12	383	343	251	197	121	100
13	634	582	430	361	195	163
14	785	851	572	615	281	287
15	1,259	1,247	954	971	470	496
16	1,431 <sup>a</sup>	1,584	1,112	1,236	620	692
17	1,785	1,766	1,425	1,355	780	784
18	2,332	2,237	1,796	1,740	1,047	1,063
19	2,359	2,242	1,713	1,678	1,082	1,021
20	2,575	2,613	1,673	1,822	1,031	1,129
21	2,502	2,473	1,602	1,527	983	939
22	2,430	2,596	1,457	1,567	915	912
23	2,419	2,410	1,316	1,340	813	829
24	2,464	2,512	1,251	1,255	778	739
25	2,457	2,450	1,182	1,139	696	720
26-29	10,420	10,047	4,239	3,997	2,463	2,530
30-34	11,429	11,635	3,952	3,802	2,566	2,150
35-39	10,409	10,744	2,746	2,874	1,570	1,611
40-44	11,764	11,273	2,637	2,373	1,418	1,297
45-49	13,418	12,366	2,799	2,409	1,599	1,438
50-54	13,365	14,278	2,570	2,505	1,589	1,538
55-59	10,555	10,607	1,774	1,794	782	1,127
60-64	6,501 <sup>a</sup>	7,573	775	1,075	434	495
65 or Older	6,257	6,647	681	655	414	394

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.11B Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2010 and 2011**

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
<b>TOTAL</b>	47.3	47.0	15.3	14.9	8.9	8.7
12	10.0	9.2	6.6	5.3	3.2	2.7
13	15.5	14.0	10.5 <sup>a</sup>	8.7	4.8	3.9
14	20.2	20.1	14.7	14.5	7.3	6.8
15	30.3	29.4	23.0	22.9	11.3	11.7
16	34.4	36.3	26.7	28.3	14.9	15.9
17	42.0	41.8	33.6	32.1	18.4	18.6
18	49.4	47.2	38.0	36.7	22.2	22.4
19	54.1	52.1	39.3	39.0	24.8	23.7
20	56.7	58.3	36.8 <sup>a</sup>	40.6	22.7	25.2
21	58.9	56.6	37.7	35.0	23.2	21.5
22	58.8	60.3	35.3	36.4	22.2	21.2
23	59.9	60.5	32.6	33.6	20.1	20.8
24	60.4	60.2	30.7	30.1	19.1	17.7
25	62.2	62.0	29.9	28.8	17.6	18.2
26-29	62.5	59.0	25.4	23.5	14.8	14.9
30-34	57.5	60.0	19.9	19.6	12.9	11.1
35-39	53.9	54.5	14.2	14.6	8.1	8.2
40-44	57.4	55.4	12.9	11.7	6.9	6.4
45-49	60.4	57.3	12.6	11.2	7.2	6.7
50-54	60.8	61.9	11.7	10.9	7.2	6.7
55-59	54.8	56.0	9.2	9.5	4.1	6.0
60-64	39.5	41.9	4.7	5.9	2.6	2.7
65 or Older	16.1	16.5	1.7	1.6	1.1	1.0

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.12A Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2010 and 2011**

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
<b>TOTAL</b>	106,613	107,842	29,301	29,739	17,409	18,071
12	39	48	35	38	10	15
13	192	213	163	168	61	88
14	412	445	356	372	167	192
15	841	890	707	757	357	378
16	1,127 <sup>a</sup>	1,279	910 <sup>a</sup>	1,033	517	594
17	1,545	1,502	1,243	1,186	691	700
18	2,029	2,018	1,564	1,581	916	965
19	2,087	2,042	1,481	1,490	947	937
20	2,277	2,392	1,456	1,638	915	1,028
21	2,281	2,264	1,364	1,349	824	861
22	2,201	2,388	1,250	1,357	800	795
23	2,228	2,198	1,130	1,141	692	701
24	2,219	2,261	1,037	1,050	664	626
25	2,206	2,233	933	948	554	613
26-29	9,468	9,063	3,340	3,280	1,901	2,096
30-34	10,269	10,442	3,082	2,760	1,956	1,625
35-39	9,100	9,503	1,894	1,994	1,127	1,234
40-44	10,661	10,065	1,702	1,606	1,017	914
45-49	12,567 <sup>a</sup>	11,364	1,881	1,641	1,120	1,038
50-54	12,414	13,204	1,598	1,815	1,072	1,129
55-59	9,674	9,797	1,250	1,317	598	855
60-64	5,988 <sup>a</sup>	7,000	542	802	316	392
65 or Older	4,787	5,230	385	416	187	295

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.12B Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2010 and 2011**

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
<b>TOTAL</b>	42.0	41.9	11.6	11.5	6.9	7.0
12	1.0	1.3	0.9	1.0	0.3	0.4
13	4.7	5.1	4.0	4.0	1.5	2.1
14	10.6	10.5	9.2	8.8	4.3	4.5
15	20.3	21.0	17.0	17.8	8.6	8.9
16	27.1	29.3	21.9	23.7	12.4	13.6
17	36.4	35.6	29.3	28.1	16.3	16.6
18	43.0	42.6	33.1	33.4	19.4	20.4
19	47.9	47.5	34.0	34.7	21.7	21.8
20	50.1	53.3	32.0 <sup>a</sup>	36.5	20.1	22.9
21	53.7	51.9	32.1	30.9	19.4	19.7
22	53.3	55.5	30.3	31.5	19.4	18.5
23	55.2	55.1	28.0	28.6	17.1	17.6
24	54.4	54.2	25.4	25.2	16.3	15.0
25	55.9	56.5	23.6	24.0	14.0	15.5
26-29	56.8	53.3	20.0	19.3	11.4	12.3
30-34	51.7	53.8	15.5	14.2	9.8	8.4
35-39	47.1	48.2	9.8	10.1	5.8	6.3
40-44	52.0	49.5	8.3	7.9	5.0	4.5
45-49	56.5 <sup>a</sup>	52.7	8.5	7.6	5.0	4.8
50-54	56.5	57.3	7.3	7.9	4.9	4.9
55-59	50.2	51.8	6.5	7.0	3.1	4.5
60-64	36.4	38.7	3.3	4.4	1.9	2.2
65 or Older	12.3	13.0	1.0	1.0	0.5	0.7

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.13A Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2010 and 2011**

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
<b>TOTAL</b>	37,361	36,921	4,533 <sup>a</sup>	3,857	1,472	1,369
12	1	4	0	1	*	*
13	10	12	2 <sup>a</sup>	9	*	1
14	22	22	14	17	1	1
15	73	76	57	46	15	21
16	101	89	71	61	22	17
17	157	135	95	95	22	28
18	287	259	157	147	47	55
19	373	360	207	200	59	62
20	509	501	200	261	57	91
21	614	527	238	227	94	68
22	685	570	252	189	64	48
23	706	645	168	169	63	45
24	687	634	184	183	53	58
25	707	756	186	188	66	37
26-29	3,177	2,933	658 <sup>b</sup>	359	157	148
30-34	3,792	3,429	644	482	246	134
35-39	3,147	3,202	350	300	116	154
40-44	4,535	3,983	329	251	96	79
45-49	6,325 <sup>b</sup>	5,239	324	314	154	136
50-54	5,479 <sup>a</sup>	6,652	231	196	112	95
55-59	3,630	3,781	104	77	4	53
60-64	1,476 <sup>a</sup>	2,073	59	77	24	34
65 or Older	870	1,040	*	8	*	5

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.13B Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2010 and 2011**

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
<b>TOTAL</b>	14.7	14.3	1.8 <sup>b</sup>	1.5	0.6	0.5
12	0.0	0.1	0.0	0.0	*	*
13	0.2	0.3	0.1 <sup>a</sup>	0.2	*	0.0
14	0.6	0.5	0.4	0.4	0.0	0.0
15	1.8	1.8	1.4	1.1	0.4	0.5
16	2.4	2.0	1.7	1.4	0.5	0.4
17	3.7	3.2	2.2	2.3	0.5	0.7
18	6.1	5.5	3.3	3.1	1.0	1.2
19	8.6	8.4	4.7	4.7	1.4	1.4
20	11.2	11.2	4.4	5.8	1.2	2.0
21	14.5	12.1	5.6	5.2	2.2	1.5
22	16.6 <sup>a</sup>	13.2	6.1	4.4	1.6	1.1
23	17.5	16.2	4.2	4.2	1.6	1.1
24	16.8	15.2	4.5	4.4	1.3	1.4
25	17.9	19.1	4.7	4.7	1.7	0.9
26-29	19.1	17.2	4.0 <sup>b</sup>	2.1	0.9	0.9
30-34	19.1	17.7	3.2	2.5	1.2	0.7
35-39	16.3	16.2	1.8	1.5	0.6	0.8
40-44	22.1 <sup>a</sup>	19.6	1.6	1.2	0.5	0.4
45-49	28.5 <sup>b</sup>	24.3	1.5	1.5	0.7	0.6
50-54	24.9	28.8	1.0	0.9	0.5	0.4
55-59	18.8	20.0	0.5	0.4	0.0 <sup>a</sup>	0.3
60-64	9.0	11.5	0.4	0.4	0.1	0.2
65 or Older	2.2	2.6	*	0.0	*	0.0

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.14A Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2010 and 2011**

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
<b>TOTAL</b>	9,208 <sup>a</sup>	8,214	885 <sup>a</sup>	625	378 <sup>a</sup>	228
12	1	1	0	1	*	*
13	5	*	0	*	*	*
14	6	5	1	3	*	*
15	9	27	4	4	2	2
16	7	10	4	9	0	0
17	32	24	9	12	1	1
18	51 <sup>a</sup>	20	26	7	10	3
19	57	47	31	16	14	3
20	87	86	8	20	*	6
21	118 <sup>b</sup>	62	17	11	7	2
22	141	99	32	16	16 <sup>a</sup>	2
23	152	117	23	11	16	2
24	131	139	19	12	5	1
25	149	143	24	14	5	2
26-29	762	635	96	36	13	7
30-34	1,107	867	37	44	4	12
35-39	973	917	113	71	37	19
40-44	1,306	1,052	132	107	68	26
45-49	1,847 <sup>b</sup>	1,214	167	123	91	84
50-54	1,248	1,551	90	51	73	37
55-59	650	702	4	23	4	14
60-64	294	397	46	29	13	*
65 or Older	76	98	*	5	*	5

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.14B Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2010 and 2011**

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
<b>TOTAL</b>	3.6 <sup>a</sup>	3.2	0.3 <sup>a</sup>	0.2	0.1 <sup>a</sup>	0.1
12	0.0	0.0	0.0	0.0	*	*
13	0.1	*	0.0	*	*	*
14	0.1	0.1	0.0	0.1	*	*
15	0.2	0.6	0.1	0.1	0.0	0.0
16	0.2	0.2	0.1	0.2	0.0	0.0
17	0.8	0.6	0.2	0.3	0.0	0.0
18	1.1 <sup>a</sup>	0.4	0.6	0.1	0.2	0.1
19	1.3	1.1	0.7	0.4	0.3	0.1
20	1.9	1.9	0.2	0.4	*	0.1
21	2.8 <sup>b</sup>	1.4	0.4	0.3	0.2	0.0
22	3.4 <sup>a</sup>	2.3	0.8	0.4	0.4 <sup>a</sup>	0.0
23	3.8	2.9	0.6	0.3	0.4	0.0
24	3.2	3.3	0.5	0.3	0.1	0.0
25	3.8	3.6	0.6	0.3	0.1	0.1
26-29	4.6	3.7	0.6	0.2	0.1	0.0
30-34	5.6	4.5	0.2	0.2	0.0	0.1
35-39	5.0	4.6	0.6	0.4	0.2	0.1
40-44	6.4	5.2	0.6	0.5	0.3	0.1
45-49	8.3 <sup>b</sup>	5.6	0.8	0.6	0.4	0.4
50-54	5.7	6.7	0.4	0.2	0.3	0.2
55-59	3.4	3.7	0.0	0.1	0.0	0.1
60-64	1.8	2.2	0.3	0.2	0.1	*
65 or Older	0.2	0.2	*	0.0	*	0.0

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.



**Table 1.15A Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2010 and 2011**

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
<b>TOTAL</b>	37,544	36,362	4,543 <sup>a</sup>	4,069	1,207 <sup>a</sup>	972
12	17	14	8	5	0	1
13	50	46	34	18	16	6
14	91	75	74	55	35	22
15	181	170	151	129	53	47
16	275	297	200	227	39 <sup>b</sup>	84
17	387	327	281 <sup>a</sup>	214	72	55
18	541	525	362	347	116	82
19	637	596	372	353	115	99
20	767	750	347	378	86	74
21	845	809	355	303	109	93
22	840	809	359	303	74	67
23	906	877	262	239	61	43
24	874	812	227	195	71	50
25	886	852	155	204	40	49
26-29	3,930	3,437	508	353	146	64
30-34	4,746	4,447	440	288	97	82
35-39	3,538	3,411	117	216	9	24
40-44	3,655	3,409	76	106	29	7
45-49	4,388	3,841	71	82	21	19
50-54	4,366	4,898	118	38	17	*
55-59	3,520	3,240	*	9	*	2
60-64	1,601	1,911	16	*	*	*
65 or Older	504	807	10	5	*	*

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.15B Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2010 and 2011**

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
<b>TOTAL</b>	14.8 <sup>a</sup>	14.1	1.8 <sup>a</sup>	1.6	0.5 <sup>a</sup>	0.4
12	0.4	0.4	0.2	0.1	0.0	0.0
13	1.2	1.1	0.8	0.4	0.4	0.1
14	2.3	1.8	1.9	1.3	0.9	0.5
15	4.4	4.0	3.6	3.0	1.3	1.1
16	6.6	6.8	4.8	5.2	0.9 <sup>a</sup>	1.9
17	9.1	7.7	6.6 <sup>a</sup>	5.1	1.7	1.3
18	11.5	11.1	7.7	7.3	2.4	1.7
19	14.6	13.9	8.5	8.2	2.6	2.3
20	16.9	16.7	7.6	8.4	1.9	1.7
21	19.9	18.5	8.4	6.9	2.6	2.1
22	20.3	18.8	8.7	7.0	1.8	1.6
23	22.4	22.0	6.5	6.0	1.5	1.1
24	21.4	19.5	5.6	4.7	1.7	1.2
25	22.4	21.6	3.9	5.2	1.0	1.2
26-29	23.6 <sup>a</sup>	20.2	3.0	2.1	0.9	0.4
30-34	23.9	22.9	2.2	1.5	0.5	0.4
35-39	18.3	17.3	0.6	1.1	0.0	0.1
40-44	17.8	16.8	0.4	0.5	0.1	0.0
45-49	19.7	17.8	0.3	0.4	0.1	0.1
50-54	19.9	21.2	0.5	0.2	0.1	*
55-59	18.3	17.1	*	0.0	*	0.0
60-64	9.7	10.6	0.1	*	*	*
65 or Older	1.3	2.0	0.0	0.0	*	*

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.16A Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2010 and 2011**

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
<b>TOTAL</b>	21,778	20,523	2,048	1,861	690	624
12	241	197	119	89	41	35
13	354	312	193	152	72	47
14	336	351	176	181	52	48
15	416	348	180	145	48	28
16	347	346	121	141	26	28
17	322	323	96	112	23	36
18	387	335	132	103	25	11
19	378	316	99	90	16	38
20	426	389	78	86	11	27
21	459 <sup>a</sup>	357	77	65	19	26
22	467	443	75	69	23	18
23	443	445	66	42	21	17
24	387	399	46	33	14	5
25	449	422	24	13	6	3
26-29	2,008	1,829	84	105	26	29
30-34	2,833 <sup>b</sup>	2,176	124	89	54	59
35-39	2,387	2,232	61	50	29	14
40-44	2,251	2,227	69	67	32	42
45-49	2,664	2,235	38	45	33	45
50-54	2,094	2,049	150	116	99	15
55-59	1,333	1,702	3	30	3	30
60-64	527	566	18	24	18	10
65 or Older	269	525	19	14	*	14

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.16B Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2010 and 2011**

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
<b>TOTAL</b>	8.6 <sup>a</sup>	8.0	0.8	0.7	0.3	0.2
12	6.3	5.3	3.1	2.4	1.1	0.9
13	8.7	7.5	4.7	3.7	1.8	1.1
14	8.6	8.3	4.5	4.3	1.3	1.1
15	10.0 <sup>a</sup>	8.2	4.3	3.4	1.2	0.7
16	8.3	7.9	2.9	3.2	0.6	0.6
17	7.6	7.7	2.3	2.7	0.6	0.8
18	8.2	7.1	2.8	2.2	0.5	0.2
19	8.7	7.3	2.3	2.1	0.4	0.9
20	9.4	8.7	1.7	1.9	0.2	0.6
21	10.8 <sup>b</sup>	8.2	1.8	1.5	0.5	0.6
22	11.3	10.3	1.8	1.6	0.5	0.4
23	11.0	11.2	1.6	1.0	0.5	0.4
24	9.5	9.6	1.1	0.8	0.3	0.1
25	11.4	10.7	0.6	0.3	0.2	0.1
26-29	12.1	10.7	0.5	0.6	0.2	0.2
30-34	14.3 <sup>b</sup>	11.2	0.6	0.5	0.3	0.3
35-39	12.4	11.3	0.3	0.3	0.1	0.1
40-44	11.0	10.9	0.3	0.3	0.2	0.2
45-49	12.0	10.4	0.2	0.2	0.1	0.2
50-54	9.5	8.9	0.7	0.5	0.4	0.1
55-59	6.9	9.0	0.0	0.2	0.0	0.2
60-64	3.2	3.1	0.1	0.1	0.1	0.1
65 or Older	0.7	1.3	0.0	0.0	*	0.0

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.17A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2010 and 2011**

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
<b>TOTAL</b>	51,832	51,243	16,051 <sup>a</sup>	14,657	6,957 <sup>a</sup>	6,119
12	180	162	119	105	75	52
13	240	213	171	129	81	52
14	293	301	200	214	98	90
15	525	480	367	356	140	134
16	603	631	437	471	162	177
17	749	735	518	475	165	183
18	1,034	922	670 <sup>a</sup>	538	277	211
19	1,097 <sup>a</sup>	941	661	574	264	230
20	1,239	1,225	660	667	268	253
21	1,290	1,210	689 <sup>a</sup>	563	286 <sup>b</sup>	193
22	1,280	1,252	640	582	263	230
23	1,230	1,269	505	501	209	196
24	1,301	1,290	528	513	218	219
25	1,361	1,264	523	427	213	178
26-29	5,155	5,030	1,792	1,569	738	693
30-34	5,426	5,364	1,666	1,687	849	745
35-39	4,492	4,259	1,175	1,147	502	484
40-44	4,668	4,489	1,134	1,176	486	459
45-49	5,526	5,092	1,100	940	502	402
50-54	5,665	6,067	1,228	919	609	506
55-59	4,317	4,581	659	599	200	250
60-64	1,975	2,485	245	270	119	103
65 or Older	2,186	1,981	365	235	233	80

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.17B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2010 and 2011**

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
<b>TOTAL</b>	20.4	19.9	6.3 <sup>b</sup>	5.7	2.7 <sup>b</sup>	2.4
12	4.7	4.3	3.1	2.8	2.0	1.4
13	5.9	5.1	4.2 <sup>a</sup>	3.1	2.0 <sup>a</sup>	1.3
14	7.5	7.1	5.2	5.1	2.5	2.1
15	12.7	11.3	8.8	8.4	3.4	3.2
16	14.5	14.4	10.5	10.8	3.9	4.1
17	17.6	17.4	12.2	11.3	3.9	4.3
18	21.9	19.5	14.2 <sup>a</sup>	11.4	5.9	4.4
19	25.1 <sup>a</sup>	21.9	15.2	13.4	6.0	5.4
20	27.3	27.3	14.5	14.9	5.9	5.6
21	30.4	27.7	16.2 <sup>b</sup>	12.9	6.7 <sup>b</sup>	4.4
22	31.0	29.1	15.5	13.5	6.4	5.3
23	30.5	31.8	12.5	12.6	5.2	4.9
24	31.9	30.9	12.9	12.3	5.3	5.3
25	34.5	32.0	13.3 <sup>a</sup>	10.8	5.4	4.5
26-29	30.9	29.6	10.8	9.2	4.4	4.1
30-34	27.3	27.7	8.4	8.7	4.3	3.8
35-39	23.3	21.6	6.1	5.8	2.6	2.5
40-44	22.8	22.1	5.5	5.8	2.4	2.3
45-49	24.9	23.6	4.9	4.4	2.3	1.9
50-54	25.8	26.3	5.6	4.0	2.8	2.2
55-59	22.4	24.2	3.4	3.2	1.0	1.3
60-64	12.0	13.7	1.5	1.5	0.7	0.6
65 or Older	5.6	4.9	0.9	0.6	0.6 <sup>a</sup>	0.2

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.18A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2010 and 2011**

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
<b>TOTAL</b>	34,908	34,247	12,242 <sup>a</sup>	11,143	5,093 <sup>a</sup>	4,471
12	155	134	108	92	71	48
13	205	177	149	111	67	42
14	244	258	160	169	83	75
15	462	398	312	297	119	116
16	539	561	374	413	125	150
17	638	621	417	401	142	132
18	864	771	525	438	220 <sup>a</sup>	149
19	947 <sup>a</sup>	785	530	444	186	177
20	1,042	987	512	492	201	164
21	1,042	974	514	416	204	147
22	1,032	1,026	506	442	208	174
23	1,022	995	371	384	153	133
24	1,065	1,025	424	403	164	150
25	1,128	1,036	397	334	174	126
26-29	4,093	4,018	1,471	1,180	544	538
30-34	4,135	3,893	1,361	1,355	626	546
35-39	3,361	2,959	924	859	422	338
40-44	3,100	2,947	796	842	346	330
45-49	3,131	3,064	826	655	395	310
50-54	2,753	3,210	762	579	367	303
55-59	1,929	2,254	452	426	104	165
60-64	1,151	1,052	178	211	67	92
65 or Older	868	1,099	171	199	103	65

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.18B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2010 and 2011**

Age Category	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
<b>TOTAL</b>	13.8	13.3	4.8 <sup>b</sup>	4.3	2.0 <sup>a</sup>	1.7
12	4.1	3.6	2.8	2.5	1.9	1.3
13	5.0	4.2	3.7	2.7	1.6	1.0
14	6.3	6.1	4.1	4.0	2.1	1.8
15	11.1 <sup>a</sup>	9.4	7.5	7.0	2.9	2.7
16	13.0	12.9	9.0	9.5	3.0	3.4
17	15.0	14.7	9.8	9.5	3.4	3.1
18	18.3	16.3	11.1	9.2	4.7 <sup>a</sup>	3.1
19	21.7 <sup>a</sup>	18.3	12.2	10.3	4.3	4.1
20	22.9	22.0	11.3	11.0	4.4	3.7
21	24.5	22.3	12.1 <sup>a</sup>	9.5	4.8 <sup>a</sup>	3.4
22	25.0	23.8	12.3	10.3	5.0	4.0
23	25.3	25.0	9.2	9.6	3.8	3.3
24	26.1	24.6	10.4	9.7	4.0	3.6
25	28.6	26.2	10.1	8.4	4.4	3.2
26-29	24.6	23.6	8.8 <sup>a</sup>	6.9	3.3	3.2
30-34	20.8	20.1	6.9	7.0	3.2	2.8
35-39	17.4	15.0	4.8	4.4	2.2	1.7
40-44	15.1	14.5	3.9	4.1	1.7	1.6
45-49	14.1	14.2	3.7	3.0	1.8	1.4
50-54	12.5	13.9	3.5	2.5	1.7	1.3
55-59	10.0	11.9	2.3	2.2	0.5	0.9
60-64	7.0	5.8	1.1	1.2	0.4	0.5
65 or Older	2.2	2.7	0.4	0.5	0.3	0.2

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.